

# Methylmercury hotspots and cycling across a High Arctic freshwater sub-catchment

VARTY, Stephanie<sup>1</sup>; LEHNHERR, Igor<sup>2</sup>; ST. PIERRE, Kyra<sup>2</sup>; KIRK, Jane<sup>3</sup>; WISNIEWSKI, Victoria<sup>1</sup>


<sup>1</sup>Department of Geography, University of Toronto, Mississauga, ON, Canada

<sup>2</sup>Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada

<sup>3</sup> Canada Centre for Inland Waters, Environment and Climate Change Canada, Burlington, ON, Canada

## Introduction

- Methylmercury (MeHg) is a toxin that bioaccumulates and biomagnifies through food webs<sup>1</sup>
- Some Arctic freshwater ecosystems produce MeHg, with most production occurring *in situ* as a result of microbial activity in sediments<sup>2,3</sup>
- There is a lack of understanding of the fate of MeHg as it is transported downstream<sup>4,5</sup> and how it varies seasonally<sup>6</sup>
- Arctic climate change has potential to alter MeHg cycling in these systems





### Objectives:

To identify where MeHg is being produced and degraded along a freshwater sub-catchment and to identify how these processes vary seasonally

Figure 1. Mercury cycling in Arctic aquatic systems

## Study Area

- A freshwater sub-catchment of Lake Hazen, Quttinirpaaq National Park, Ellesmere Island, Nunavut
- The Skeleton Lake continuum allows for samples to be taken as water travels into Lake Hazen passing through a lake, two wetland ponds, a wetland and a tundra creek channel



## Sampling Schematic



## Methods

### Using Stable Hg isotope tracers to investigate methylation potential:

- Incubations of lake/pond water, lake/pond sediments, wetland soils, snowpack and snow melt samples were conducted shortly before melt, during the ice-covered period during and the ice-free, productive period to explore the fate of MeHg as it is transported through the Skeleton Lake sub-catchment
- Inorganic <sup>198</sup>Hg and Me<sup>199</sup>Hg were injected into sample media at predetermined concentrations to track short-term methylation and demethylation potential, respectively<sup>9,10</sup>




Figure 2. Illustrations of stable Hg isotope injection methods

### Spatiotemporal Hg & MeHg Survey:

- Wetland soil cores were taken in summer 2016 along multiple transects perpendicular to water flow to examine saturation gradient
- Snow & snow melt samples were taken in spring 2017
- %MeHg obtained from soils, water, snow and melt will be used as a proxy for longer-term methylation potential<sup>11</sup>



## Results – High levels of MeHg in Spring




Figure 3. Total MeHg concentrations (ngL<sup>-1</sup>) in Skeleton Lake water during ice-covered and ice-free period




Figure 4. MeHg concentrations (ngL<sup>-1</sup>) in snowpack along Skeleton Continuum, May 2017

## Results – Spatial Variation




Figure 5. MeHg variation in water throughout the Skeleton Lake Sub-catchment in 2016




Figure 6. Potential methylation in Skeleton Continuum, Summer 2016

Figure 7. Potential demethylation in Skeleton Continuum, Summer 2016

## Discussion

- Preliminary spring results show that snow (0.53 ngL<sup>-1</sup>) and Sk. Lake bottom water (0.56ngL<sup>-1</sup>) have elevated MeHg concentrations before/during melt
- Spring methylation experiments in the water column/snow will shed light on observed seasonal patterns
- Skeleton Lake and Pond 11 sediments both act as hotspots for methylation, providing MeHg to the downstream wetland
- Potential demethylation in wetland soils is significantly higher when compared pond/lake sediments, establishing that there is potential for the wetland to act as a sink of MeHg through demethylation in soils
- Additionally, MeHg in the water flowing into the wetland is largely particulate bound, suggesting that sorption of MeHg onto soil organic matter may play a role in decreased MeHg concentrations<sup>4</sup>
- System MeHg & DOC concentrations have a significant positive regression (p=0.03, R<sup>2</sup>=0.22)

## Future Steps

- Mechanisms driving changes in %MeHg can be elucidated by exploring the fate of MeHg as it is transported downstream
- Analysis of snowpack, snowmelt and snow particulate matter will provide an understanding of the origin of MeHg in snowpack and how it is transported and transformed during accumulation and melt
- Comparison of ice-free and ice-covered periods will aid in identifying how climate change may alter MeHg cycling in the Arctic

## Acknowledgements

- Vincent St. Louis, Maria Cavaco, Jess Serbu, Graham Colby, Charlie Talbot, Allie Ho and Amber Gleason
- ArcticNet, NSERC, Northern Contaminants Program, Polar Continental Shelf Program

Stephanie Varty, MSc Candidate  
University of Toronto, Dept. of Geography  
Email: stephanie.varty@mail.utoronto.ca

Igor Lehnerr, Assistant Professor  
University of Toronto, Dept. of Geography  
Email: igor.lehnerr@utoronto.ca

## References

- [1] Lehnerr, et al. 2012a. *Environmental Science Technology*, 46:10523-10531. [2] Lehnerr et al. 2012b. *Environmental Science Technology*, 46:10523-10531. [3] Hammerschmidt et al. 2006. *Environmental Science Technology*, 40:1204-1211. [4] Lossetto et al. 2004. *Environmental Science Technology* 38:3004-3010. [5] Olfert & Sicilliano. 2009. *Science of the Total Environment*, 407:5169-1700. [6] Steffen. *Environment Canada, Clean Air Regulatory Agenda Mercury Science Program*, in press [7] ESRI 2011. ArcGIS Desktop: Release 10. Redlands, CA: Environmental Systems Research Institute. [8] Government of Canada: Natural Resources Canada; Earth Sciences Sector. 2016. CanVec [9] Mao et al. 2011. *Applied Mechanics and Materials*, 71-78: 3201-3206 [10] Lehnerr, et al. 2011. *Nature Geoscience*, 4(5), 298-302. [11] Eckley & Hintelmann. 2006. *Science of the Total Environment*, 368, 111-125.