

Introduction

- Future oil production (Fig. 1) combined with growing vessel traffic¹ (Fig. 2) increase the risk of an oil spill event off the coast of Labrador.
- Sea ice and iceberg floes may impede conventional oil spill response measures².




Figure 1.

The Labrador Shelf is the focus of ongoing oil and gas exploration via one of the world's largest seismic survey programs.

Seismic lines shown in blue and red along the Labrador Shelf. Red box indicating parcels up for bid in 2018.

Maps modified from TGS (www.tgs.com) and CNLOPB (www.cnlopb.ca)




Figure 2.

As Canada's Northwest Passage becomes increasingly ice free, shipping routes are predicted to increase along the Labrador coast¹ (★).

- Bioremediation of spilled oil by indigenous hydrocarbonoclastic bacteria (HCB)³ is essential for eliminating oil that enters marine systems⁴.
- Biostimulation, supplementation of nutrients (e.g., nitrogen and phosphorous) to stimulate HCBs, has proven to be effective for nearshore marine oil spills⁵.
- Whether indigenous microbial communities can effectively biodegrade spilled hydrocarbons off the coast of Labrador remains poorly understood.
- **The purpose of this research is to probe the coast of Labrador for a microbial response to hydrocarbon contamination, focusing on biostimulation as an approach to spilled oil in cold, ice impacted marine environments.**

Hypotheses

1. Hydrocarbon degrading microbes are present in Labrador Shelf sediments and poised to respond in the event of an oil spill
2. Indigenous microbes are capable of degrading hydrocarbons aerobically at 4°C
3. Biostimulation with nitrogen and phosphorous will increase the rate of biodegradation

Objectives

1. Uncover baseline microbial community composition of Labrador Shelf sediment
2. Demonstrate response of indigenous microbes to biostimulation in mock oil spill microcosms incubated at aerobically at 4°C
3. Assess temporal shifts in microbial communities during nutrient and hydrocarbon amended experiments

Materials & Methods




Figure 3. Sediment obtained from Labrador Shelf⁶ (Station 640⁶) aboard the CCGS Amundsen⁶ in 2015 by box corer⁶




Figure 4. Microcosm components. Four treatment conditions vary by diesel and nutrient concentrations.




Figure 5. In-house Illumina MiSeq used for 16S rRNA gene sequencing of amplicons prepared with 731F and 806R primers.



Figure 6. Microbial respiration (CO₂ and O₂) in microcosm headspace monitored by GC as indicator of hydrocarbon degradation. GC-MS used to compare hydrocarbon profiles before and after microbial degradation incubations.

Figure 7. Bioinformatic analysis using in-house MetaAmp⁷ pipeline.

Results




Figure 8. Microbial respiration, indicated by O₂ consumption and CO₂ production, in sediment microcosms incubated at 4°C and amended with either 0.1 or 1% v/v diesel fuel, with either low or high concentrations of N and P (either 15 or 4670 μM for P, and either 2 or 1470 μM for N). Mean values from triplicate incubations are plotted, with error bars showing standard deviations. High nutrient with 1% diesel treatments had five times the CO₂ production, with full oxygen consumption after 71 days of incubation compared to the other three treatment conditions.




Figure 9. Bacterial community composition revealed by 16S rRNA gene amplicon libraries prepared from Labrador Shelf sediment microcosms amended with high levels of nutrient (4670 μM: P 1470 μM: N) and 1% (v/v) diesel. Triplicate samples were assessed at day 0 and day 29 of a 71-day incubation at 4°C. Incubations showed increases in the relative abundance of Proteobacteria, specifically genera known to include HCBs: *Oleispira*, *Thalassolituus*, *Perlicidibaca*, and *Neptunomonas*.

Summary & Future Work

- Genera containing HCBs are present in Labrador Shelf sediment and shown to increase in relative abundance when incubated with diesel and high nutrients.
- Biodegradation in the environment with ambient nutrient concentrations may be slow without supplementation of nitrogen and phosphorous.
- Future work involves hydrocarbon profiling with GC-MS to observe degradation before and after incubation, as well as repeating this experiment using crude oil.

References

1. Smith, L. C., and Stephenson, S. R. (2013). New Trans-Arctic shipping routes navigable by midcentury. *Proceedings of the National Academy of Sciences*, 110(13), E1191-E1195.
2. Fisheries and Oceans Canada (2009). Environmental Response [Brochure]. Ottawa, Ontario: Fisheries and Oceans Canada, Canadian Coast Guard Maritime Services.
3. Berry, D., and Gutierrez, T. (2017). Evaluating the detection of hydrocarbon-degrading bacteria in 16S rRNA gene sequencing surveys. *Frontiers in microbiology*, 8(896).
4. Head, I. M., Jones, D. M., and Røling, W. F. (2006). Marine microorganisms make a meal of oil. *Nature Reviews Microbiology*, 4(3), 368(6470), 413-418.
5. Bragg, J. R., Prince, R. C., Harner, E. J., and Atlas, R. M. (1994). Effectiveness of bioremediation for the Exxon Valdez oil spill. *Nature*, 368(6470), 413-418.
6. Google Maps. (2017). Labrador Shelf [Satellite image]. Retrieved from URL: www.google.ca/maps/@54.1428768,-62.1830727,2685206m/data=!3m1!1e3
7. Dong, X., Kleiner, M., Sharp, C. E., Thorson, E., Li, C., Liu, D., and Strous, M. (2017). Fast and simple analysis of MiSeq amplicon sequencing data with MetaAmp. *Frontiers in Microbiology*, 8(1461). doi: 10.3389/fmicb.2017.01461

